

CHAPTER 35

EARLY EFFECTS OF RADIATION

Early Effects of Radiation

- A radiation response in human within a few days to months
- It is described as *deterministic*

Deterministic Radiation Response

- Biologic response whose severity varies with radiation dose
- A dose threshold usually exists

ACUTE RADIATION LETHALITY

Death

- The most devastating human response to radiation exposure

Acute Radiation-Induced Lethality

- It is of only academic interest in diagnostic radiology

PRINCIPAL EARLY EFFECTS OF RADIATION EXPOSURE ON HUMANS & THE APPROXIMATE THRESHOLD DOSE

Effect	Anatomic Site	Threshold Dose
Death	Whole body	200 rad/2 Gy _t
Hematologic depression	Whole body	25 rad/250 mGy _t
Skin erythema	Small field	200 rad/2 Gy _t
Epilation	Small field	300 rad/3 Gy _t
Chromosome aberration	Whole body	5 rad/50 mGy _t
Gonadal dysfunction	Local tissue	10 rad/100 mGy _t

Diagnostic x-ray beams always result in partial-body exposure, which is less harmful than whole-body exposure!

Chernobyl Incident

- April 1986

Three Mile Island Incident

- March 1979

Acute Radiation Syndrome

- Radiation sickness that occurs in human after the whole-body dose s of 1 Gy (100 rad) or more of ionizing radiation delivered over a short time

Three Syndromes

- Hematologic Death, Gastrointestinal (GI) Death & Central Nervous System (CNS) Death

Prodromal Period

- The immediate response of radiation sickness
- *Approximate Dose:* > 100 rad
- *Mean Survival Time:* —
- *Clinical S&S:* nausea, vomiting & diarrhea

Latent Period

- The time after exposure during which there is no sign of radiation sickness
- *Approximate Dose:* 100-10,000 rad
- *Mean Survival Time:* —
- *Clinical S&S:* none

Hematologic Syndrome

- It is characterized by a reduction in white cells, red cells & platelets
- *Approximate Dose:* 200-1000 rad
- *Mean Survival Time:* 10-60 days
- *Clinical S&S:* nausea, vomiting, diarrhea, anemia, leukopenia, hemorrhage, fever & infection
- *Prodromal Period:* mild symptoms
- *Latent Period:* general feeling of wellness
- *Period of Manifest Illness:* vomiting, mild diarrhea, malaise, lethargy & fever
- *Recovery:* 2-4 weeks or 6 months (full)
- *Cause of Death:* generalized infection, electrolyte imbalance & dehydration

GI Period

- It occurs principally because of severe damage to the cells lining the intestines
- *Approximate Dose:* 1000-5000 rad
- *Mean Survival Time:* 4-10 days

CHAPTER 35

EARLY EFFECTS OF RADIATION

- *Clinical S&S*: same as hematologic plus electrolyte imbalance, lethargy, fatigue & shock
- *Prodromal Period*: vomiting & diarrhea
- *Latent Period*: no symptoms present
- *Period of Manifest Illness*: second wave of nausea & vomiting, followed by diarrhea, anorexia
- *Cause of Death*: unprevented rapid progression of symptoms

CNS Period

- Its ultimate cause is elevated fluid content of the brain
- *Characterized By*: increased intracranial pressure, vasculitis & meningitis
- *Approximate Dose*: > 5000 rad
- *Mean Survival Time*: 0-3 days
- *Clinical S&S*: same as GI plus ataxia, edema, system vasculitis & meningitis
- *Prodromal Period*: severe nausea & vomiting
- *Latent Period*: earlier symptoms disappear
- *Period of Manifest Illness*: more severe prodromal symptoms, disoriented, loss muscle coordination, dyspnea, convulsive seizures, loss of equilibrium, ataxia & lethargy

LD_{50/60}

- The dose of radiation to the whole body that causes 50% of irradiated subjects to die within 60 days
- It quantitatively measured the acute radiation lethality
- *Humans*: 350 rad

Acute radiation lethality follows a nonlinear, threshold dose-response relationship!

Mean Survival Time

- Average time between exposure & death
- *Hematologic Syndrome*: dose dependent
- *GI Syndrome*: remain constant
- *CNS Syndrome*: dose dependent

LOCAL TISSUE DAMAGE

Local Tissue Damage

- It follows a threshold-type dose response relationship
- *Characteristic*: deterministic response

Local Tissues That Can Be Affected Immediately

- Skin
- Gonads
- Bone marrow

Partial-Body Irradiation

- A higher dose is required to produce a response
- It affects organ & tissue
- *Effect*: cell death
 - *Result*: shrinkage of the organ or tissue

Atrophy

- The shrinkage of an organ or tissue due to cell death

EFFECT ON SKIN

Skin

- The tissue with which we have had the most experience
- Three Layers
 - *Epidermis*: outer layer
 - *Basal Cells*: its lowest layer
 - *Dermis*: intermediate layer of connective tissue
 - *Subcutaneous*: layer of fat & connective tissue
- *Other Accessory Structures*: hair follicles, sweat glands & sensory receptors
- *Cells Replacement Rate*: 2 %/day (50 % for GI)
- *Skin Effects*: nonlinear, threshold dose-response relationship

Basal Cells

- The stem cells that mature as they migrate to the surface of the epidermis

CHAPTER 35

EARLY EFFECTS OF RADIATION

Damage to basal cells results in the earliest manifestation of radiation injury to the skin!

Orthovoltage X-rays

- Range: 200-300 kVp

Erythema

- A sunburn-like reddening of the skin
- The first observed biologic response to radiation exposure

Desquamation

- Ulceration & denudation of the skin

Moist Desquamation

- The clinical intolerance for radiation therapy

X-ray-Induced Erythema

- One of the hazards to the patient the early of radiology

Skin-Erythema Dose (SED)

- Dose of radiation, usually about 200 rad, that causes redness of the skin

Epilation/Alopecia/Fox Mange

- Loss of hair

SED₅₀

- The dose required to affect 50% of those irradiated
- Dose: 500 rad

POTENTIAL RADIATION RESPONSES OF SKIN FROM HIGH-DOSE FLUOROSCOPY

Potential Radiation Response	Threshold Dose	Approximate Time of Onset
Early transient erythema	200 rad	Hours
Main erythema	600 rad	10 days
Temporary epilation	300 rad	3 weeks
Permanent epilation	700 rad	3 weeks
Moist desquamation	1500 rad	4 weeks

Grenz Rays

- It is used to treat tinea capitis (ringworm)
- Range: 10-20 kVp

EFFECTS ON GONADS

Testes

- The male gonads
- It produces *spermatogonia* & matures into *sperm*

Ovaries

- The female gonads
- It produces *oogonia* & matures into *ovum*

Germ Cells

- Produced by both ovaries & testes

Gametogenesis

- The process of development of germ cells

Progression of Germ Cell

- *Male*: Spermatogonia (most radiosensitive) → Spermatocyte → Spermatid → Sperm
- *Female*: Primordial Follicle → Mature Follicle (most radiosensitive) → Corpus Luteum → Ovum

Oogonia

- The stem cells of the ovaries
- They multiply in number only before birth & during fetal life

Primordial Follicles

- They grow to encapsulate the oogonia

Oocyte

- A matured oogonia

Ovum

- A mature female germ cell
- *Fertilization*: 400-500 ova
 - Number of years of menstruation times 13 per year

CHAPTER 35

EARLY EFFECTS OF RADIATION

Spermatogonia

- The stem cells of the testes
- Continually being produced from stem cells progressively through a number of stage to maturity

Spermatocyte

- A matured spermatogonia

Spermatid

- A matured spermatocyte

Spermatozoa/Sperm

- A mature male germ cell
- *Maturation Process: 3-5 weeks*

Ovaries

- Irradiation Causes
 - *Early Life:* atrophy
 - *After Puberty:* suppression & delay of menstruation
- *10 rad:* suppresses menstruation
- *25-50 rad:* increased genetic mutations
- *200 rad:* temporary sterility
- *500 rad:* sterility

The most radiosensitive cell during female germ cell development is the oocyte in the mature follicle!

Testes

- *Irradiation Causes:* atrophy
- *10 rad:* reduce the number of spermatozoa
- *200 rad:* temporary sterility
- *500 rad:* sterility

Spermatogonial Stem Cells

- The most sensitive phase in the gametogenesis of the spermatozoa

Male Gametogenesis

- A self-renewing system

HEMATOLOGIC EFFECTS

Periodic Blood Examination

- The only monitoring performed on x-ray & radium workers before
- Total cell counts & a white cell differential count

Under no circumstances is a periodic blood examination recommended as a feature of any current radiation protection program!

Hematologic Depression

- *Threshold Dose: 25 rad*

Hemopoietic System

- Another example of cell renewal system
 - Same with gametogenesis
- Bone marrow
- Circulating blood
- Lymphoid tissue
 - Lymph nodes, Spleen & Thymus
- *Principal Effect of Radiation:*
 - Depressed number of blood cells in the peripheral circulation

Pluripotential Stem Cell

- Stem cell that has the ability to develop into several different types of mature cells
- It produces lymphocytes, granulocytes, thrombocytes & erythrocyte

Lymphocytes/White Blood Cells

- Blood cells involved in the immune response
- Manufactured by spleen & thymus
- *Lifetime in the Bone Marrow:* varying (hours or years)
- *Lifetime in the Peripheral Blood:* varying (hours or years)

Granulocytes

- Scavenger type of cells used to fight bacteria
- *Lifetime in the Bone Marrow:* 8-10 days
- *Lifetime in the Peripheral Blood:* couple of days

CHAPTER 35

EARLY EFFECTS OF RADIATION

- *Recovery: 2 months*

Thrombocytes/Platelets

- Blood cells involved in the clotting of blood to prevent hemorrhage
- *Lifetime in the Bone Marrow: 5 days*
- *Lifetime in the Peripheral Blood: 1 week*
- *Recovery: 2 months*

Erythrocytes/Red Blood Cells

- Blood cells that are transportation agents for oxygen
- Less sensitive than the other blood cells
 - *Rationale: long lifetime in the peripheral blood*
- *Lifetime in the Bone Marrow: 8-10 days*
- *Lifetime in the Peripheral Blood: 4 months*
- *Recovery: 6 months to 1 year*

Bone Marrow

- It manufactures most circulating blood cells including lymphocytes
- *Child: uniformly distributed throughout the skeleton*
- *Adult: restricted to flat bones such as ribs, sternum, skull & ends of long bones*

HEMOPOIETIC CELL SURVIVAL

Principal Radiation Response of Hemopoietic System

- Decrease in the number of all types of blood cells in the circulating peripheral blood

Lethal Injury

- It causes depletion of mature circulating cells

Lymphopenia

- Reduced in number of lymphocytes

The lymphocytes & the spermatogonia are the most radiosensitive cells in the body!

Granulocytosis

- Rapid rise in number of granulocytes

Granucytopenia

- Rapid decrease & slower decrease of granulocytes

Thrombocytopenia

- Depletion of platelets

CYTOGENETIC EFFECTS

Cytogenetics

- The study of the genetics of cells particularly cell chromosomes

Radiation-induced chromosome aberrations follow a nonthreshold dose-response relationship!

Human Peripheral Lymphocytes

- Most often used for cytogenetic analysis

Karyotype

- A chromosome map

Each cell consists of 22 pairs of autosomes & a pair of sex chromosomes – the X-chromosomes from the female & the Y chromosomes from the male!

Chromosomes Structural Radiation Damage

- Single-Hit Chromosome Aberrations
- Double-Hit Chromosome Aberrations

Reciprocal Translocation

- It requires a karyotype for detection

Point Genetic Mutations

- Undetectable even with karyotype construction

Hit

- Radiation interaction with chromosomes

DNA Hit

- It results in an invisible disruption of the molecular structure of the DNA

CHAPTER 35

EARLY EFFECTS OF RADIATION

Chromosome Hit

- It produces a visible derangement of the chromosome
- It represents severe damage to the DNA

Singe-Hit Chromosome Aberration

- Visualized & recorded during the M phase
- Irradiation During G₁ Phase
 - *Cause:* chromatid break
 - *During S Phase:* replicated
 - *During Metaphase:* a chromosome with material missing from the ends of two sister chromatids & two acentric fragments
- Irradiation During G₂ Phase
 - *Causes:* single or double chromatid break
 - *During S Phase:* replicated
 - *During Metaphase:* a chromosome with an arm that is obviously missing genetic material & a chromatid fragment

Chromatid Deletion

- The breakage of a chromatid

Isochromatid

- Fragments in chromosome aberrations

Acentric

- Without a centromere

Multi-Hit Chromosome Aberration

- Not uncommon
- Irradiation During G₁ Phase
 - *Causes:* ring & dicentric chromosomes
 - *Ring:* when two hits occur on the same chromosome
 - *Dicentric:* when adjacent chromosome each suffer one hit & recombine
- *Irradiation During G₂ Phase:* similar to G₁ phase but rarer

Stickiness

- A condition in which the mechanism for the joining of chromatids depends

Radiation-Induced Reciprocal Translocation

- Multi-hit chromosome that require karyotype analysis for detection
- *Results in:*
 - No loss of genetic material
 - Simply a rearrangement of the genes

Kinetics of Chromosome Aberration

- *Single-Hit Aberration:*
 - It occurs at very low doses of radiation
 - *Dose-Response Relationship:* linear, nonthreshold
- *Multi-Hit Aberration:*
 - It occurs when the radiation dose exceeds approximately 100 rad
 - *Dose-Response Relationship:* nonlinear, nonthreshold

Radiation Dose-Response Relationship For Cytogenetic Damage

- *Single-Hit:* $Y = a + bD$
- *Multi-Hit:* $Y = a + bD + cD^2$

Cytogenetic Analysis

- Biologic radiation dosimeter

Approximate Chromosome Aberration Frequency

- Two single-hit aberrations per rad per 1000 cells
- One multi-hit aberration per 10 rad per 1000 cells